

Efficacy of prostaglandin on ovary regarding reproduction in the fish, *Cyprinus carpio* (L.)

Veena Mahajan

Assistant Professor & Head, Department of Zoology, J. M. Patel College Bhandara, Maharashtra, India

E-mail: mahajanveena2020@gmail.com

Manuscript Details

Received :13.05.2020

Accepted: 29.06.2021

Published: 30.06.2021

Available online on <https://www.irjse.in>

ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Veena Mahajan. Efficacy of prostaglandin on ovary regarding reproduction in the fish, *Cyprinus carpio* (L.), *Int. Res. Journal of Science & Engineering*, 2021, Volume 9(3): 91-94.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Abstract

Prostaglandin plays an important role in ovulation, luteal function, implantation and maintenance of gestation. It is a class of fatty acids which are traditionally associated with variety of autocrine and paracrine functions in vertebrate body. Ovary has a definite role in the reproductive physiology. Prostaglandin was injected 1.5ml/kg body weight at every alternate days for one week intramuscularly in the fish, *Cyprinus carpio*. It has been observed that due to prostaglandin, oogenesis stimulated in the ovary which was mediated through the gonadotrophic hormone. Thus, it had been concluded that prostaglandin is an inducer of spawning reflexes in the fish, *Cyprinus carpio* that induces gonadal maturity and used to achieve early breeding.

Keywords: Autocrine, paracrine, prostaglandin, reproductive physiology, spawning reflexes.

1. Introduction

Prostaglandins were first discovered and isolated from human semen in 1930s by Ulf Von Euler and colleagues of Sweden. Prostaglandin is a wonder biomolecule derived from 20 c atoms polyunsaturated fatty acids (PUFA), mainly released from membrane phospholipid like phosphatidylcholine or choline phosphoglycerides. They are basically derived from prostanoic acid, a 20 c fatty acid with cyclopentene ring. Prostaglandin F reaches a maximum in ovulated follicle at the time of ovulation [1]. Prostaglandin also improves sperm quantity in the canine ejaculate would benefit

all assisted reproductive techniques used in all species [1]. In fish, exogenous prostaglandins also show stimulatory effect on the induction of ovulation. Stacey [2] showed that indomethacin blocks ovulation induced by HCG injection in goldfish, but exogenous prostaglandin overcome the indomethacin blockade. Jalabert [3] observed that ovulation of trout follicles matured in vivo could be induced in vitro by adding prostaglandin F_{2α}. The effects of prostaglandin on the spawning behavior of female goldfish were also observed by Stacey, [2]. Recently, existence and role of prostaglandin in fishes have been studied by several authors in many fishes. F prostaglandins also function as a hormone that stimulates female sexual behavior [4, 5, 6]. At the time of ovulation female goldfish produce large quantities of prostaglandin (PGF₂) which acts as hormone to trigger spawning (oviposition) behaviors [7, 8] before being metabolized and released where males recognized it as a releaser pheromone [9, 10, 11]. Female sexual behavior apparently is synchronized with ovulation by mechanisms which respond to elevated plasma prostaglandin as an indicator of the presence of ovulated eggs [7]. Prostaglandin plays important role in ovulation, luteal function, implantation and maintenance of gestation. They are used to synchronise estrus and induce parturition. F prostaglandin rise at the time of ovulation and travel to the brain where they elicit female sexual behavior. Thus, the effect of prostaglandin on ovary in relation to reproduction in the fish, *Cyprinus carpio* (L.) is meagre. Moreover, such studies are limited to foreign species and hence, the study is undertaken on these aspects in locally available carp. The present work is an attempt in this direction to assess the suitability of prostaglandin as an inducer for breeding purpose.

2. Materials and Method

Cyprinus carpio were collected, reared in a fibre glass tanks and acclimatized to the laboratory conditions. The

experiment was carried out for one week. Control and experimental groups were formed. Fishes from control groups were injected with 1.5ml of distilled water and experimental groups were injected with 1.5ml of prostaglandin intramuscularly at every alternate day for one week. At the end of the experiment, fishes from control and experimental groups were sacrificed to study the histomorphological changes in the ovary.

3. Results

CONTROL GROUP:

At the end of the experiment, the survival rate of control fish was 100%. The average weight length and girth of female was 610gm, 28cm and 26cm. The ovary was in the maturing form and histologically, reorganized into young oocytes, early maturing oocytes, advanced maturing oocytes and few mature oocytes (prespawning oocyte). The ovaries were yellowish in colour. Histologically oocyte showed small clear yolk vesicles and nucleus with undulated nuclear membrane.

EXPERIMENTAL GROUP:

At the end of the experiment, the survival rate of the fish was 100%. The average weight, length and girth of female were 620gm, 28cm and 27cm. At the end of the experiment, the ovaries were in prespawning phase. The entire body cavity was occupied by the ovaries which were turgid and deep yellow in colour. The number of ova could be seen by the naked eye through the thin ovarian wall. The ova were opaque as well as translucent. The fish was gravid with rounded abdomen. On pressing, eggs oozed out. At the end of the experiment, histologically, a large number of oocytes with fused yolk vesicles, yolk globules and migrating nucleus along with ripe eggs were seen in a section and became ready to spawn. A fully ripe gravid female has soft bulging rounded abdomen with swollen reddish vent.




Figure 1 A: Section of ovary showing oocytes at different stages of development including developing oocytes, primary oocyte, maturing oocytes, yolk vesicles yolk granules, and atretic oocytes X10.
B: maturing oocytes with nucleus showing outer zona granulosa and inner zona radiata Iron haematoxylin X10.
C: Mature oocyte showing interstitial cells, Mallory's triple strain X10.
D: Section of ovary showing different stages of developing oocytes Iron haematoxylin

4. Discussion

The exact biochemical mechanism of ovulation in fish has not been fully elucidated. Very little information is available on ovarian maturation in fishes [12]. It is indicated that corticosteroids derived from the ovary may act as a local hormone mediating pituitary gonadotropins induce ovulation in *Orizias latipus*, [12] but in rainbow trout, northern pike and gold fish 17α hydroxy- 20β -dihydroprogesterone plays a preferential role in initiating maturation and ovulation may be triggered in vitro by prostaglandin $F_2\alpha$ action on mature follicles. The present results reveal that prostaglandin $F_2\alpha$ plays an important role in maturation and ovulation of fish *Cyprinus carpio*. Male and female both exhibit an increased frequency of courtship behavior following prostaglandin $F_2\alpha$ treatment which is also supported by the views of Cole and Smith [13]. They also reported that prostaglandin $F_2\alpha$ appears to induce the release of a female specific chemical in fathead minnows that triggers courting behavior in conspecific male. Present study investigated the possibility of prostaglandin $F_2\alpha$ may have pheromonal role in *Cyprinus carpio* as it

stimulates both male and female to breed easily by exhibiting vigorous courtship. In the presence study, $PGF_2\alpha$ also stimulates oocyte maturation in *Cyprinus carpio*. At the time of ovulation female gold fish oviducts synchronise and secrete $PGF_2\alpha$ that induces reproductive behaviour [8;10]. $PGF_2\alpha$ and its metabolites mainly 15 keto - $PGF_2\alpha$ are also released into water as post ovulatory pheromone that induce male spawning behavior and further increase male GTH - II and sperm production to stimulate the GTH - II [4, 9,10]. Present studies also indicate that F prostaglandin is metabolised and released to the water where it functions as a sex pheromone. Therefore, $PGF_2\alpha$ also plays a dual role as a para hormone and hormone, synchronising male and female sexual behavior in *Cyprinus carpio*. Davidson *et al.* [5] have found that exposure to water borne $PGF_2\alpha$ increased neurogenesis and GnRH concentration in male goldfish brain and modulate brain plasticity associated with behavioral changes during spawning season via the neuroendocrine (GnRH) and motor components of the pheromone-reproductive system. This finding confirms the above view in *Cyprinus carpio* in relation with

increased number of gonadotropins, olfactory secretion, increased secretory activity in thyroid and adrenal gland. GSI and fecundity is also increased significantly.

5. Summary and Conclusion

The objective of the present study was to evaluate the effect of prostaglandin on ovary in relation to reproduction in the fish, *Cyprinus carpio*. The effect of prostaglandin on ovary showed stimulatory effect. Particularly, the gonads (ovary) were excessively stimulated resulted into early maturity. Thus, prostaglandin had stimulatory effect on the gonadal secretion.

Thus, the results obtained from the present study, it can be concluded that utility prostaglandin in the fish breeding technology will be effective as it stimulates the oogenesis in the ovary of *Cyprinus carpio* which is mediated through the gonadotrophic hormone resulted in to the secretion of estrogen which is essential for the sexual behavior in the fish breeding and prostaglandin used as the stimulant in the fish breeding. Prostaglandin also used to achieve early maturity.

Conflict of interest

No conflict of interest influenced in this research.

6. References

1. Milan B Hess. The effect of prostaglandin F_{2α}, oxytocin and gonadotropin releasing hormone in ejaculate characteristic in the Dog. Virginia polytechnic institute , 2002; (thesis).
2. Stacey NE. Effect of indomethacin and prostaglandins on the spawning behavior of female goldfish. Prostaglandin 12: 113 - 126.ish. Gen. Comp. endocrinal. 1976; 100, 128 - 142.
3. Jalabert, B. In vitro oocyte maturation and ovulation in rainbow trout, *Salmo gairdneri*, Northern pike, *Essox lucius* and *Carassius auratus*. J. Fish. Res. Board can, 1976; .33: 9.
4. Sorensen PW and Goetz FW. Pheromonal and reproductive functions of F prostaglandin and their metabolites in teleost fish. J. Lipid Med. 1993; 6: 385 - 393.
5. Davidson, Yu - Wen - Chung. C. B. Rees., Mara. B. Bryan and W. Li. Neurogenic and neuroendocrine effects of goldfish pheromones. The Journal of neuroscience, 2008 ; Dec. 28 (53): 14492 - 14499.
6. Fedrica Di Costanzo et al., Prostaglandin in marine organisms: A review. Mar. Drugs, 2019; 17, 428, doi103390/mar070428
7. Stacey NE. Hormones, pheromones and reproductive behavior. J. Fish. Physiol. / Biochem 2003; 28: 229 - 235. (Vol. 28: No. 1 - 4).
8. Stacey NE and Peter RE. Central action of prostaglandin in spawning behavior of female goldfish. Physiol Behav. 1979; 22: 1191 - 1196.
9. Sorensen PW, Hara TJ, Stacey NE and Goetz FW. F-Prostaglandin functions as potent olfactory stimulants that comprise the post - ovulatory female sex pheromone in goldfish. Biol. Reprod., 1983 ; 39: 1039 - 1050. : 217 - 32.g. Virginia.4.
10. Sorensen PW and Stacey NE and Chamberlain KJ. Differing behavioral and endocrinological effects of two female sex pheromones on male goldfish. Horn. Behav, 1989; 23 (3)
11. Stacey NE and Sorensen PW. Fish hormonal Pheromones. In: Hormones, brain and behavior, Vol. 2 (Palfalvi, Arnold, D., Ategen. A, Farbachs Rubin R. eds), 2002; Pp. 3375 - 435. New York: Academy.
12. Hirose K. Endocrine control of ovulation in medaka (*Oryzias latipes*) and Ayu (*Plecoglossus altivelis*). J. Fish. Res. 1976; Boar.
13. Cole KS and Smith RJ. Release of chemicals by prostaglandin-treated female Fathead minnows, *Pimephales promelas* that stimulate male courtship. Horn Behav. 1987; (4): 440 - 56.