

Diversity of Algal Flora in Salekasa Tehsil of Gondia District, Maharashtra, India

Nandeshwar MP, Mendhe BK and Pardhi BN

Shankaralal Agrawal Science College Salekasa (M.S.), India

Email: mpnandeshwar@gmail.com

Manuscript Details

Available online on <https://www.irjse.in>
ISSN: 2322-0015

Cite this article as:

Nandeshwar MP, Mendhe BK and Pardhi BN. Diversity of Algal Flora in Salekasa Tehsil of Gondia District, Maharashtra., *Int. Res. Journal of Science & Engineering*, 2023, Special Issue A13:75-80.
<https://doi.org/10.5281/zenodo.XXXXXX>

Article published in Special issue of National Conference on "New Frontier of Biological Sciences (NCNFBS-2023) jointly organized by Internal Quality Assurance Cell (IQAC) and Biological Society, Shri. Shivaii Education Society Amravati's Science College, Pawni, Dist. Bhandara, Maharashtra, India, date, April 26, 2023.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

ABSTRACT

Salekasa is a tribal Tehsil in the District of Gondia, Maharashtra, India. Situated at the *Latitude is 21.30286 and Longitude are 80.56443*. Geographically it is in the east part of the Gondia District and it shared its border with Dongargarh Tahsil of Chhattisgarh State. More than 50 species were found in this study but only 50 species were studied and recorded in this article. It contains algae belonging to *Cyanophyceae, Chlorophyceae, Charophyceae, Euglenophyceae, and Bacillariophyceae*. All 50 species were found growing luxuriantly in different seasons. Out of these all 20 species of *Cyanophyceae*, 19 species of *Chlorophyceae*, 2 species of *Charophyceae*, 4 species of *Euglenophyceae*, and 5 species of *Bacillariophyceae*.

Keywords: Freshwater lakes, Diversity of Algae, Salekasa Tehsil.

Introduction

Water is most important for living organisms, purification of water today needs that and it makes use for the living organism. Many anthropogenic activities are responsible for the contamination of water and then it results in the degradation of aquatic ecosystems. The water bodies contain many biological components. Macrofloral diversity is protected from environmental degradation. Aquatic flora plays a key role to maintain the food chain and physico-chemical parameters of the water bodies. Hence throughout the world, many researchers have worked on aquatic flora. Algal community encountered in the water body reflects the average ecological condition and therefore they may be used as an indicator of water quality [1, 2]

Seasonal phytoplanktonic diversity of Kitham Lake, [3]. The Study on Phytoplankton in Kandhar, Nanded District by [4]. Algal flora of Navegaon bandh, Gondia district [5]. A qualitative and quantitative study of phytoplankton of River Wainganga near Markandadeo, Dist. Gadchiroli (M.S.) by Tijare [6]. Assessment of aquatic macrophytes diversity from Karmaveer Kannamwar (Dina Project) reservoir Regadi of Chamorshi tehsil district Gadchiroli [7]. New records of freshwater algae for Maharashtra state: Investigation from the major rivers of Chandrapur district by Reddy [8].

Methodology

Algal flora samples were collected at monthly intervals from June 2021 to October 2022. Various floating, submerged, and attached epiphytic algal samples were collected from the selected site namely Salekasa Lake, Pangaon Lake, Halbitola Lake, Rondha Nalha, Paol Dawana Lake, Kahali Lake, Sonpuri Lake, Tirkhedi Lake, etc. Google map photos are in Fig. 1. Collected samples were washed with an acid-washed bottle and preserved in 4% formalin. After investigation identification of algal flora with the help of Monographs and standard literature [9].




Figure : 1. Salekasa lake, 2. Pangaon Lake, 3. Halbitola Lake, 4. Rondha Nalha, 5. Paul Dawana Lake, 6. Kahali Lake, 7. Tirkhedi lake, and 8. Sonpuri Lake (Courtesy-Google map photos).

Result & Discussion

The water quality of freshwater bodies mostly depends on the composition of flora and fauna and some abiotic factors. Eutrophication in freshwater bodies is an indicator of the presence of algal bloom of cyanobacteria. Also, the high-density population of Cyanobacteria in the lake is harmful to the other organism of that ecosystem because they produce certain hepatotoxic and neurotoxic substances. In the present study, 50 species were studied. It contains algae belonging to *Cynophyceae* (Blue-green Algae), *Chlorophyceae* (Green Algae), *Charophyceae* (Stoneworts), *Euglenophyceae* (Euglenoids), and *Bacillariophyceae* (Diatoms). All 50 species were found growing luxuriantly in different seasons. Out of these all 20

species of *Cyanophyceae*, 19 species of *Chlorophyceae*, 2 species of *Charophyceae*, 4 species of *Euglenophyceae*, and 5 species of *Bacillariophyceae*.




Fig. 2: Distribution of recorded species in the families

Table 1: Algal Diversity

| Sr.No. | Name of Algae | Rainy Session | Winter Session | Summer Session |
|--|--|---------------|----------------|----------------|
| Family:- <i>Cynophyceae</i> (Blue-Green Algae) – 20 Species | | | | |
| 1 | <i>Aphanocapsa grevillei</i> (Hass.) Rabenh. | - | - | + |
| 2 | <i>Aulosira fertilissima</i> (Rio De Janeiro) | - | + | - |
| 3 | <i>Gloeocapsa nigrescens</i> Nag. | - | + | - |
| 4 | <i>Microcystis flos-aquae</i> (Wittrok Kirchner 1898) | - | - | + |
| 5 | <i>Microcystis marginata</i> (Meneghini, Kuetzing, 1846) | - | + | - |
| 6 | <i>Merismopedia glauca</i> forma (Rao, C. S.) | - | - | + |
| 7 | <i>Merismopedia punctata</i> Meyen. | + | + | + |
| 8 | <i>Spirulina gigantea</i> Schmidle | - | + | - |
| 9 | <i>Oscillatoria chalybea</i> v. <i>insularis</i> Gardner | - | + | + |
| 10 | <i>Oscillatoria proboscidea</i> Gom. (after Gomont) | - | + | + |
| 11 | <i>Oscillatoria subbrevis</i> Schmidle | + | + | - |
| 12 | <i>Phormidium tenue</i> (Menegh.) Gom. (after Fremy). | - | + | + |
| 13 | <i>Lyngbya digueti</i> Gomont. | + | + | + |
| 14 | <i>Lyngbya hieronmusii</i> Lemm. | - | + | - |
| 15 | <i>Lyngbya majuscula</i> Havery ex. Gomont. | - | - | + |
| 16 | <i>Nostoc calcicola</i> Breb. (after Fremy) | + | + | + |
| 17 | <i>Nostoc piscinale</i> Kutz. (after Fremy) | - | + | + |
| 18 | <i>Anabaena circunalis</i> Rabenhorst ex Born. et. Flah. | - | + | + |
| 19 | <i>Scytonema cincinnatum</i> Thuret (after Fremy) | - | + | + |
| 20 | <i>Scytonema javanicum</i> (Kutz.) Bornet | + | + | + |

Table 1: Continued...

| Sr.No. | Name of Algae | Rainy Session | Winter Session | Summer Session |
|--|---|---------------|----------------|----------------|
| Family:- Chlorophyceae (Green Algae) - 19 Species | | | | |
| 21 | <i>Chlorococcum humicola</i> (Naegeli) Rabenhorst | + | + | + |
| 22 | <i>Chroococcus minutus</i> (Kutz. Nageli 1849) | + | + | + |
| 23 | <i>Chroococcus tenax</i> (Kirchn.) Hieron | + | + | + |
| 24 | <i>Pediastrum duplex</i> Meyen var. <i>asperum</i> | + | + | + |
| 25 | <i>Pediastrum simplex</i> v. <i>duodenarium</i> (Bail.) | - | + | - |
| 26 | <i>Pediastrum tetras</i> (Ehr.) Ralfs. | + | + | + |
| 27 | <i>Tetraedron trigonum</i> (Naeg.) | - | + | - |
| 28 | <i>Scenedesmus dimorphus</i> Lemm | + | - | - |
| 29 | <i>Scenedesmus arcuatus</i> Lemm | + | - | - |
| 30 | <i>Scenedesmus quadricauda</i> v. <i>longispina</i> | - | + | - |
| 31 | <i>Oedogonium globosum</i> Nordstedt ex Hirn. | - | + | - |
| 32 | <i>Spirogyra ellipsospora</i> Transeau 1914. | + | + | + |
| 33 | <i>Spirogyra hyalina</i> Cleve (Transeau f) | + | - | + |
| 34 | <i>Spirogyra mirabilis</i> (Hassall) Kuetzing | + | - | + |
| 35 | <i>Closterium cyanthia</i> DeNot | + | + | - |
| 36 | <i>Closterium didymotocum</i> Corda. | + | + | - |
| 37 | <i>Closterium ehrenbergii</i> Menegh. | - | + | + |
| 38 | <i>Closterium moniliferum</i> (Bory) Ehr. | + | - | + |
| 39 | <i>Cosmarium contractum</i> Kirchner. | + | + | + |
| Family:- Charophyceae (Stoneworts) – 02 Species | | | | |
| 40 | <i>Chara</i> Sp. | + | + | + |
| 41 | <i>Nitella</i> Sp. | + | + | + |
| Family:- Euglenophyceae (Euglenoids) – 04 Species | | | | |
| 42 | <i>Euglena acus</i> Ehrenberg (Gojdics f) | - | + | + |
| 43 | <i>Euglena mutabilis</i> Schmitz. (Gojdics) | - | + | + |
| 44 | <i>Phacus acuminatus</i> Stokes. Hueb. | - | + | + |
| 45 | <i>Phacus longicaudus</i> (Her.) Duj. | + | + | + |
| Family:- Bacillariophyceae (Diatoms) – 05 Species | | | | |
| 46 | <i>Gyrosigma baikalensis</i> Skv. | + | - | - |
| 47 | <i>Gyrosigma maharashtrensis</i> sp. Nov. | + | + | - |
| 48 | <i>Navicula halophila</i> (Grun.) Cleve f. <i>robusta</i> | - | + | + |
| 49 | <i>Navicula microcephala</i> Grun | - | + | + |
| 50 | <i>Nitzschia irremissa</i> Cholnoky | + | + | + |
| Chart for. all 50 species were found growing luxuriantly in different seasons. Out of these all 20 species of <i>Cyanophyceae</i> , 19 species of <i>Chlorophyceae</i> , 2 species of <i>Charophyceae</i> , 4 species of <i>Euglenophyceae</i> , and 5 species of <i>Bacillariophyceae</i> | | | | |




Fig. 1.3: Algal Photos arranged in alphabetically

Conclusion

Conflicts of interest: The authors stated that no conflicts of interest.

References

1. Bhatt LR, Lecoul P, Lekhal HD, Jha PK. Physico-chemical characteristics and phytoplankton for Taudha Lake, Kathmandu. *Poll.Res.*, 1999, 18 (4): 353-358.
2. Saha SB, Bhattacharya SB, Choudhary. A Diversity of phytoplankton of sewage pollution brackish water tidal ecosystems. *Environ.Biol.*, 2000, 21 (1): 9-14.
3. Tiwari A and Chauhan SV. Seasonal phytoplanktonic diversity of Kitham Lake, Agra. *J Environ Biol.*, 2006, 27: 35-38.
4. Pawar SK, Pulle JS and Shendge KM. The study on phytoplankton of pethwadaj Dam, TalulaKandhar, District - Nanded, Maharashtra, *J. Aqua. Biol*, 2006, 21 (1): 1-6.
5. Shahare PC. Algal flora of Navegaon bandh, Gondia district. (Maharashtra). *I J R B A T*, May 2014, Issue (2), Volume-II, e-ISSN 2347 - 517X
6. Tijare RV. Qualitative and quantitative study of phytoplankton of River Wainganga near Markandadeo, Dist. Gadchiroli (M.S.) *Environment Conservation Journal*, 2020, 21 (3): 43-49, 2020 ISSN 0972-3099 (Print) 2278-5124 (Online)
7. Meshram MP, Tijare RV, Zod R. Assessment of aquatic macrophytes diversity from Karmaveer Kannamwar (Dina Project) reservoir Regadi of Chamorshi tehsil district Gadchiroli November 2020, DOI:[10.32381/IPSJ.2020.36.1-2.9](https://doi.org/10.32381/IPSJ.2020.36.1-2.9)
8. B. Mallesh Raj Reddy. New records of freshwater algae for Maharashtra state: Investigation from the major rivers of Chandrapur district: *An NeBIO.*, 2021, Vol. 12, No. 2, Jun 2021.
9. Desikachary TV. Cyanophyta ICAR monographs on Algae New Delhi.1959, PP 686.

Publisher's Note

IRJSE remains neutral with regard to jurisdictional claims in published maps and institutional affiliations

Submit your manuscript to a IJLSCI journal and benefit from:

- ✓ Convenient online submission
- ✓ Rigorous peer review
- ✓ Immediate publication on acceptance
- ✓ Open access: articles freely available online
- ✓ High visibility within the field

Submit your next manuscript to IRJSE through our manuscript management system uploading at the menu "**Make a Submission**" on journal website

Email your next manuscript to IRJSE
editor@irjse.in